

## Parameterizing the Intersection of a Sphere and a Plane

**Problem:** Parameterize the curve of intersection of the sphere  $\mathcal{S}$  and the plane  $\mathcal{P}$  given by

$$(\mathcal{S}) \quad x^2 + y^2 + z^2 = 9$$

$$(\mathcal{P}) \quad x + y = 2$$

**Solution:** There is no foolproof method, but here is one method that works in this case and many others where we are intersecting a cylinder or sphere (or other “quadric” surface, a concept we’ll talk about Friday) with a plane.

**Step 1:** Find an equation satisfied by the points of intersection in terms of two of the coordinates.

We’ll eliminate the variable  $y$ . Note that the equation  $(\mathcal{P})$  implies  $y = 2 - x$ , and substituting this into equation  $(\mathcal{S})$  gives:

$$x^2 + (2 - x)^2 + z^2 = 9$$

$$x^2 + 4 - 4x + x^2 + z^2 = 9$$

$$2x^2 - 4x + z^2 = 5$$

**Step 2:** Parameterize the equation from Step 1, writing the two coordinates (here,  $x$  and  $z$ ) in terms of a parameter  $t$ .

There are many ways to parameterize

$$2x^2 - 4x + z^2 = 5 \tag{1}$$

but one way is to note that an equation of the form

$$u^2 + v^2 = R^2$$

for a constant  $R$  and variables  $u$  and  $v$  is the equation of a circle of radius  $R$  in  $(u, v)$ -coordinates which can be parameterized using trigonometric functions as

$$\begin{cases} u = R \cos t \\ v = R \sin t, \end{cases} \quad 0 \leq t \leq 2\pi \tag{2}$$

For equation (1), we can complete the square:

$$2x^2 - 4x + z^2 = 5$$

$$2(x^2 - 2x) + z^2 = 5$$

$$2[(x - 1)^2 - 1] + z^2 = 5$$

$$2(x - 1)^2 + z^2 = 7$$

which is of the correct form for  $u = \sqrt{2}(x - 1)$ ,  $v = z$ , and  $R = \sqrt{7}$ . The parameterization (2) becomes:

$$\begin{cases} \sqrt{2}(x - 1) = \sqrt{7} \cos t \\ z = \sqrt{7} \sin t, \end{cases} \quad 0 \leq t \leq 2\pi$$

and solving for  $x$  and  $z$  we get a parameterization for two of the three coordinates:

$$\begin{cases} x = \sqrt{\frac{7}{2}} \cos t + 1 \\ z = \sqrt{7} \sin t, \end{cases} \quad 0 \leq t \leq 2\pi$$

**Step 3:** *The final step (which is barely even a step) is to add a parameterization for the final coordinate.*

From the plane equation ( $\mathcal{P}$ ), we know  $y = 2 - x$ , so we can substitute in the parameterization for  $x$  to get:

$$y = 2 - x = 2 - \left(\sqrt{\frac{7}{2}} \cos t + 1\right) = 1 - \sqrt{\frac{7}{2}} \cos t$$

The final parameterization for all three coordinates is:

$$\begin{cases} x = \sqrt{\frac{7}{2}} \cos t + 1 \\ y = 1 - \sqrt{\frac{7}{2}} \cos t \\ z = \sqrt{7} \sin t, \end{cases} \quad 0 \leq t \leq 2\pi$$

and so

$$\mathbf{r}(t) = \left\langle \sqrt{\frac{7}{2}} \cos t + 1, 1 - \sqrt{\frac{7}{2}} \cos t, \sqrt{7} \sin t \right\rangle, \quad 0 \leq t \leq 2\pi$$