

# MATHEMATICS 263 December 2004 Final Exam Solutions

- 1) The temperature in the solid ellipsoid  $x^2 + xz + \frac{3}{2}z^2 + 2(y-2)^2 \leq 11$  is given by

$$T(x, y, z) = \sqrt{y+3} e^{2x-z}$$

- (a) Find a line that is perpendicular to the surface of the ellipsoid and passes through the point  $P = (1, 1, 2)$ . Call this line  $L$ .  
 (b) Calculate the rate of temperature change per unit distance at  $P$  in the direction inward along  $L$ .  
 (c) Estimate the temperature of the solid 0.09 units from point  $P$  inward along  $L$ .

**Solution.** (a) Let  $g(x, y, z)$  be the LHS of the inequality. Then, an (outward pointing) normal to the surface of the solid at  $(1, 1, 2)$  is given by

$$\vec{N} = \nabla g(1, 1, 2) = \langle 2x + z, 4(y - 2), x + 3z \rangle|_{(1,1,2)} = \langle 4, -4, 7 \rangle$$

Therefore, the normal line at this point can be written

$$\boxed{\vec{r}(t) = \langle 1, 1, 2 \rangle + t \langle 4, -4, 7 \rangle, \quad -\infty < t < \infty}$$

- (b) We want the directional derivative at  $P$  in the direction of the *inward*-pointing normal, that is in the direction

$$\hat{n} = \frac{-\vec{N}}{|\vec{N}|} = \frac{\langle -4, 4, -7 \rangle}{\sqrt{(-4)^2 + 4^2 + (-7)^2}} = \left\langle -\frac{4}{9}, \frac{4}{9}, -\frac{7}{9} \right\rangle$$

Since we have

$$\nabla T(x, y, z) = \left\langle 2\sqrt{y+3}e^{2x-z}, \frac{1}{2}(y+3)^{-1/2}e^{2x-z}, -\sqrt{y+3}e^{2x-z} \right\rangle$$

we may calculate the desired rate of change

$$D_{\hat{n}}T(1, 1, 2) = \hat{n} \cdot \nabla T(1, 1, 2) = \left\langle -\frac{4}{9}, \frac{4}{9}, -\frac{7}{9} \right\rangle \cdot \left\langle 4, \frac{1}{4}, -2 \right\rangle = \boxed{-\frac{1}{9}}$$

- (c) The temperature at the new point  $P'$  may be linearly approximated by

$$T(P') = T(P) + 0.09 D_{\hat{n}}T(P) = 2 + 0.09 \left(-\frac{1}{9}\right) = 2 - 0.01 = \boxed{1.99}$$

- 2) The mass  $m$  of an object with kinetic energy  $E$  and speed  $v$  is  $m = 2E/v^2$ . If a body has a measured kinetic energy of 200 and a measured speed of 100, but the measurements could have an error of  $\pm 1\%$ , what is the approximate maximum percentage error in the calculated value of the mass?

**Solution.** First note that  $\frac{\partial m}{\partial E} = \frac{2}{v^2}$  and  $\frac{\partial m}{\partial v} = -4\frac{E}{v^3}$ . Also, we know that  $\Delta E = \pm(200)(.01) = \pm 2$  and  $\Delta v = \pm(100)(.01) = \pm 1$ . We need to compute  $\Delta m$ .

$$\begin{aligned} \Delta m &= m(200 + \Delta E, 100 + \Delta v) - m(200, 100) \\ &\approx \nabla m(200, 100) \cdot \langle \Delta E, \Delta v \rangle \quad \text{by linear approximation} \\ &= \frac{\partial m}{\partial E}(200, 100)\Delta E + \frac{\partial m}{\partial v}(200, 100)\Delta v \\ &= \frac{2}{100^2}(\pm 2) - 4\frac{200}{100^3}(\pm 1) \\ &= \frac{\pm 4}{100^2} + \frac{\mp 8}{100^2} \\ &= \frac{\pm 4 \mp 8}{100^2} = \frac{\pm 12}{100^2} = \frac{\pm 12}{10,000} \end{aligned}$$

Since  $m \approx 2\frac{200}{100^2} = \frac{4}{100}$  and  $\pm\frac{12}{10,000} = \frac{4}{100}(\pm\frac{3}{100})$  the approximate error in the mass is  $\boxed{\pm 3\%}$ .

3) Find the minimum and maximum values of  $x^2 + 2y^2 - x$  in the region  $x^2 + y^2 \leq 1$ .

**Solution.** Write  $f(x, y) = x^2 + 2y^2 - x$ . If the minimum/maximum value of  $f$  is achieved in  $x^2 + y^2 < 1$ , it must be achieved at a critical point. The critical points are the solutions of

$$0 = f_x = 2x - 1 \quad 0 = f_y = 4y$$

So the only critical point is  $(\frac{1}{2}, 0)$ , where  $f$  takes the value  $-\frac{1}{4}$ . The other possibility is that  $f$  takes its min/max value on  $x^2 + y^2 = 1$ . The value of  $f(x, y)$  at  $(x, y) = (\cos \theta, \sin \theta)$  is  $g(\theta) = \cos^2 \theta + 2 \sin^2 \theta - \cos \theta$ . So the minimum/maximum value of  $f$  on the boundary is the same as the minimum/maximum value of  $g(\theta)$ , which we determine by finding the critical points of  $g(\theta)$ .

$$0 = g'(\theta) = -2 \sin \theta \cos \theta + 4 \sin \theta \cos \theta + \sin \theta = \sin \theta (2 \cos \theta + 1)$$

Hence the critical points are at

$$\sin \theta = 0 \iff y = 0 \iff (x, y) = (\pm 1, 0) \quad \cos \theta = -\frac{1}{2} \iff x = -\frac{1}{2} \iff (x, y) = (-\frac{1}{2}, \pm \frac{\sqrt{3}}{2})$$

From the table of all possible candidates, below, we see that the minimum  $\boxed{-\frac{1}{4}}$  and the maximum is  $\boxed{\frac{9}{4}}$ .

$(x, y)$	$(\frac{1}{2}, 0)$	$(1, 0)$	$(-1, 0)$	$(-\frac{1}{2}, \frac{\sqrt{3}}{2})$	$(-\frac{1}{2}, -\frac{\sqrt{3}}{2})$
$f(x, y)$	$-\frac{1}{4}$	0	2	$\frac{9}{4}$	$\frac{9}{4}$

4) Convert to polar coordinates and evaluate:

$$I = \int_0^2 \int_0^{\sqrt{2x-x^2}} (k + 3\sqrt{x^2 + y^2}) dy dx.$$

Express your answer in terms of the constant  $k$ .

**Solution.** Name the domain of integration  $D$ . This lies inside the vertical strip  $0 \leq x \leq 2$ , where it runs up from  $y = 0$  to the curve where

$$y^2 = 2x - x^2, \quad \text{i.e.,} \quad x^2 - 2x + y^2 = 0, \quad \text{i.e.,} \quad (x - 1)^2 + y^2 = 1.$$

Hence  $D$  is a semicircle of radius 1. In polar coordinates,

$$x^2 + y^2 = 2x \iff r^2 = 2r \cos \theta \iff r = 2 \cos \theta,$$

and the angles of interest obey  $0 \leq \theta \leq \pi/2$ , so

$$I = \int_{\theta=0}^{\pi/2} \int_{r=0}^{2 \cos \theta} (k + 3r)r dr d\theta = \int_{\theta=0}^{\pi/2} \left[ \frac{k}{2} r^2 + r^3 \right]_{r=0}^{2 \cos \theta} d\theta = \int_0^{\pi/2} [2k \cos^2 \theta + 8 \cos^3 \theta] d\theta$$

From the formula sheet  $\int_0^{\pi/2} \cos^2 \theta d\theta = \frac{\pi}{4}$  and  $\int_0^{\pi/2} \cos^3 \theta d\theta = \frac{2}{3}$  so that  $\boxed{I = \frac{k\pi}{2} + \frac{16}{3}}$ . Notice that the coefficient of  $k$  can be found using geometry: it's just the area of the semicircle  $D$ , which is  $\pi/2$ .

5) Let  $\vec{F}(x, y) = (\sin y + y \cos x)\hat{i} + (\sin x + x \cos y)\hat{j}$ .

(a) Determine whether or not  $\vec{F}$  is conservative. If it is, find a potential function for  $\vec{F}$ .

(b) Calculate  $\int_C \vec{F} \cdot d\vec{r}$ , where  $C$  is the piece of the parabola  $y = \frac{2}{\pi}x^2$  from  $A = (0, 0)$  to  $B = (\frac{\pi}{2}, \frac{\pi}{2})$ .

**Solution.** (a)  $\vec{F}$  might be conservative if  $\frac{\partial F_1}{\partial y} = \frac{\partial F_2}{\partial x}$ . In this case,

$$\frac{\partial F_1}{\partial y} = \frac{\partial}{\partial y}(\sin y + y \cos x) = \cos y + \cos x = \frac{\partial}{\partial x}(\sin x + x \cos y) = \frac{\partial F_2}{\partial x}$$

so we need to find a function  $\varphi(x, y)$  such that  $\nabla\varphi = \vec{F}$ . There are many ways to find such a function. We could, by inspection, guess that  $\varphi(x, y) = x \sin y + y \sin x$  would work and then verify that  $\nabla\varphi(x, y) = (\sin y + y \cos x)\hat{i} + (x \cos y + \sin x)\hat{j} = \vec{F}$ , as desired. Let's try a more mechanical way to find this function. To have  $\frac{\partial\varphi}{\partial x} = F_1$  we need

$$\varphi(x, y) = \int F_1 dx = \int (\sin y + y \cos x) dx = x \sin y + y \sin x + C(y)$$

To also have  $\frac{\partial\varphi}{\partial y} = F_2$  we need  $x \cos y + \sin x + C'(y) = F_2 = \sin x + x \cos y$ . Thus  $C'(y) = 0$ , and hence  $C(y)$  is a constant. We are free to choose  $C(y) = 0$  so that  $\varphi(x, y) = x \sin y + y \sin x$ .

(b) The specified work integral is  $\phi(B) - \phi(A) = \pi$ .

6) Let  $D$  be the solid that is bounded below by the plane  $2x + 2y + z + 2 = 0$  and is bounded above by the paraboloid  $z = 4 - (x + 1)^2 - (y + 1)^2$ . Let the field  $\vec{F}$  be given by

$$\vec{F}(x, y, z) = \frac{\langle y, 1, z \rangle}{\sqrt{x^2 + y^2}}$$

(a) Parameterize the curve of intersection of the plane and paraboloid in terms of the polar coordinate  $\theta$ .

(b) Let  $S_1$  be the portion of the surface of  $D$  formed by the paraboloid. Parameterize  $S_1$ .

(c) Let  $J$  denote the flux through  $S_1$  into the solid  $D$ . Express  $J$  as an iterated double integral using the parameterization of part (b). Evaluate the inner integral. Evaluation of the remaining outer integral is not required.

**Solution.** (a) The curve of intersection is given by the system

$$2x + 2y + z + 2 = 0 \quad z = 4 - (x + 1)^2 - (y + 1)^2$$

Substituting the value for  $z$  given by the second equation into the first gives  $2x + 2y + 4 - (x + 1)^2 - (y + 1)^2 + 2 = 0$  or  $4 - x^2 - y^2 = 0$ , the circle centered at the origin of radius 2. This yields the parameterization

$$x = 2 \cos \theta \quad y = 2 \sin \theta \quad z = -2 - 2x - 2y = -2 - 4 \cos \theta - 4 \sin \theta \quad 0 \leq \theta \leq 2\pi$$

(b) From part (a), we know that the  $(x, y)$  coordinates of the points of  $S_1$  will be contained within the disc  $x^2 + y^2 \leq 4$ , so using cylindrical coordinates, we have

$$x = r \cos \theta \quad y = r \sin \theta \quad z = 4 - (r \cos \theta + 1)^2 - (r \sin \theta + 1)^2 = 2 - r^2 - 2r \cos \theta - 2r \sin \theta \quad 0 \leq r \leq 2, 0 \leq \theta \leq 2\pi$$

(c) For the parameterization

$$\vec{r}(r, \theta) = \langle r \cos \theta, r \sin \theta, 2 - r^2 - 2r \cos \theta - 2r \sin \theta \rangle$$

we have

$$\begin{aligned}
\frac{\partial \vec{r}}{\partial r} \times \frac{\partial \vec{r}}{\partial \theta} &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \cos \theta & \sin \theta & -2r - 2 \cos \theta - 2 \sin \theta \\ -r \sin \theta & r \cos \theta & 2r \sin \theta - 2r \cos \theta \end{vmatrix} \\
&= \hat{i} \begin{vmatrix} \sin \theta & -2r - 2 \cos \theta - 2 \sin \theta \\ r \cos \theta & 2r \sin \theta - 2r \cos \theta \end{vmatrix} - \hat{j} \begin{vmatrix} \cos \theta & -2r - 2 \cos \theta - 2 \sin \theta \\ -r \sin \theta & 2r \sin \theta - 2r \cos \theta \end{vmatrix} + \hat{k} \begin{vmatrix} \cos \theta & \sin \theta \\ -r \sin \theta & r \cos \theta \end{vmatrix} \\
&= \langle 2r + 2r^2 \cos \theta, 2r + 2r^2 \sin \theta, r \rangle
\end{aligned}$$

As the third component is positive, this is an upward normal and so outward from the surface. Therefore, we want to use

$$\hat{n} dS = - \left( \frac{\partial \vec{r}}{\partial r} \times \frac{\partial \vec{r}}{\partial \theta} \right) dr d\theta = - \langle 2r + 2r^2 \cos \theta, 2r + 2r^2 \sin \theta, r \rangle dr d\theta$$

to calculate our flux integral. Converting  $\vec{F}$  to cylindrical coordinated yields

$$\vec{F} = \langle \sin \theta, 1/r, 2/r - r - 2 \cos \theta - 2 \sin \theta \rangle$$

and so our flux integral is

$$\begin{aligned}
J &= \iint_{S_1} \vec{F} \cdot \hat{n} dS = \int_0^2 dr \int_0^{2\pi} d\theta [\langle \sin \theta, \frac{1}{r}, \frac{2}{r} - r - 2 \cos \theta - 2 \sin \theta \rangle \\
&\quad \bullet (- \langle 2r + 2r^2 \cos \theta, 2r + 2r^2 \sin \theta, r \rangle)] \\
&= \boxed{\int_0^2 dr \int_0^{2\pi} d\theta [-2r \sin \theta - 2r^2 \sin \theta \cos \theta - 4 + r^2 + 2r \cos \theta]} \\
&= \int_0^2 dr \int_0^{2\pi} d\theta [-2r \sin \theta - r^2 \sin 2\theta - 4 + r^2 + 2r \cos \theta] \\
&= \int_0^2 dr [2r \cos \theta + \frac{1}{2} r^2 \cos 2\theta - 4\theta + r^2 \theta + 2r \sin \theta]_{\theta=0}^{\theta=2\pi} \\
&= \boxed{\int_0^2 dr [-8\pi + 2\pi r^2]} = -8\pi r + \frac{2}{3}\pi r^3 \Big|_{r=0}^{r=2} = -16\pi + \frac{16}{3}\pi = \boxed{-\frac{32}{3}\pi}
\end{aligned}$$

For the other order of integration

$$\begin{aligned}
J &= \int_0^{2\pi} d\theta \int_0^2 dr [-2r \sin \theta - 2r^2 \sin \theta \cos \theta - 4 + r^2 + 2r \cos \theta] \\
&= \int_0^{2\pi} d\theta [-4 \sin \theta - \frac{16}{3} \sin \theta \cos \theta - 8 + \frac{8}{3} + 4 \cos \theta] \\
&= \boxed{\int_0^{2\pi} d\theta [-4 \sin \theta - \frac{16}{3} \sin \theta \cos \theta - \frac{16}{3} + 4 \cos \theta]}
\end{aligned}$$

7) Let  $\mathcal{R}$  denote the solid region defined by the simultaneous inequalities

$$x \geq 0, \quad y \geq 0, \quad z \geq 0, \quad 1 \leq x^2 + y^2 + z^2 \leq 4$$

Let  $\mathcal{S}$  denote the surface of  $\mathcal{R}$ .

(a) Sketch  $\mathcal{R}$  and  $\mathcal{S}$ .

(b) Evaluate the outward flux of the following vector field through  $\mathcal{S}$ :

$$\vec{\mathbf{F}}(x, y, z) = \langle x^5 + y \sin(z), y^5 + z \sin(x), 10x^2 y^2 z - x \rangle.$$

(c) Find the flux of  $\vec{\mathbf{F}}$  downward through the bottom of  $\mathcal{S}$ , i.e., through the flat part of  $\mathcal{S}$  that lies in the plane  $z = 0$ .

**Solution.** (a)  $\mathcal{R}$  is one quarter of the solid between a sphere of radius 1 and a sphere of radius 2.

(b) Here

$$\nabla \cdot \vec{\mathbf{F}} = [5x^4] + [5y^4] + [10x^2 y^2] = 5(x^4 + 2x^2 y^2 + y^4) = 5(x^2 + y^2)^2.$$

By the Divergence Theorem, the desired flux is

$$J = \iint_{\mathcal{S}} \vec{\mathbf{F}} \cdot \hat{\mathbf{n}} dS = \iiint_{\mathcal{R}} \nabla \cdot \vec{\mathbf{F}} dV = 5 \iiint_{\mathcal{R}} (x^2 + y^2)^2 dV.$$

Spherical coordinates are convenient for the volume integral.

$$\begin{aligned} J &= 5 \iiint_{\mathcal{R}} (x^2 + y^2)^2 dV = 5 \int_{\theta=0}^{\pi/2} \int_{\phi=0}^{\pi/2} \int_{\rho=1}^2 [\rho^2 \sin^2 \phi]^2 \rho^2 \sin \phi d\rho d\phi d\theta \\ &= 5 \int_{\theta=0}^{\pi/2} \int_{\phi=0}^{\pi/2} \frac{1}{7} [\rho^7]_{\rho=1}^2 \sin^5 \phi d\phi d\theta = 5 \frac{127}{7} \left(\frac{\pi}{2}\right) \int_0^{\pi/2} \sin^5 \phi d\phi \end{aligned}$$

From the formula sheet  $\int_0^{\pi/2} \sin^5 \theta d\theta = \frac{8}{15}$  so that  $J = \frac{4 \times 127}{7 \times 3} \pi$ .

(c) Write  $\mathcal{S}_0$  for the bottom of  $\mathcal{S}$ . On  $\mathcal{S}_0$  we have  $z = 0$ ; the outward unit normal is  $\hat{\mathbf{n}} = -\mathbf{k}$ , so

$$\vec{\mathbf{F}} \cdot \hat{\mathbf{n}} = x.$$

Polar coordinates work well for this:

$$\begin{aligned} \iint_{\mathcal{S}_0} \vec{\mathbf{F}} \cdot \hat{\mathbf{n}} dS &= \iint_{\mathcal{S}_0} x dA = \int_{\theta=0}^{\pi/2} \int_{r=1}^2 [r \cos \theta] r dr d\theta \\ &= \left( \int_{\theta=0}^{\pi/2} \cos \theta d\theta \right) \left( \int_{r=1}^2 r^2 dr \right) = (1) \left( \frac{2^3 - 1^3}{3} \right) = \boxed{\frac{7}{3}} \end{aligned}$$

- 8) Let  $\vec{\mathbf{F}}(x, y, z) = (e^{x^2} + y)\hat{\mathbf{i}} + (\sin(y^3) + xz)\hat{\mathbf{j}} + z^2\hat{\mathbf{k}}$ . Use Stokes's theorem to evaluate  $\int_{\mathcal{C}} \vec{\mathbf{F}} \cdot d\vec{\mathbf{r}}$  where  $\mathcal{C}$  is the curve  $x^2 + y^2 = 10$ ,  $x + y + z = 4$  with positive orientation (i.e. counter-clockwise) as viewed from high on the  $z$ -axis.

**Solution.** Let  $D$  denote the disk  $x + y + z = 4$ ,  $x^2 + y^2 \leq 10$  and let  $\hat{\mathbf{n}}$  denote the upward pointing unit normal to  $D$ . By Stokes' theorem

$$\int_{\mathcal{C}} \vec{\mathbf{F}} \cdot d\vec{\mathbf{r}} = \iint_D \nabla \times \vec{\mathbf{F}} \cdot \hat{\mathbf{n}} dS$$

For the specified vector field

$$\nabla \times \vec{\mathbf{F}} = -x\hat{\mathbf{i}} + (z - 1)\hat{\mathbf{k}}$$

Viewing  $x + y + z = 4$  as  $z = f(x, y)$  with  $f(x, y) = 4 - x - y$

$$\begin{aligned} \hat{\mathbf{n}} dS &= (-f_x \hat{\mathbf{i}} - f_y \hat{\mathbf{j}} + \hat{\mathbf{k}}) dxdy = (\hat{\mathbf{i}} + \hat{\mathbf{j}} + \hat{\mathbf{k}}) dxdy \\ \nabla \times \vec{\mathbf{F}} \cdot \hat{\mathbf{n}} dS &= (-x, 0, f(x, y) - 1) \cdot (1, 1, 1) dxdy = (-x + f(x, y) - 1) dxdy \\ &= (3 - 2x - y) dxdy \end{aligned}$$

Since  $x$  and  $y$  are odd functions,

$$\int_{\mathcal{C}} \vec{\mathbf{F}} \cdot d\vec{\mathbf{r}} = \iint_{x^2+y^2 \leq 10} (3 - 2x - y) dxdy = 3 \iint_{x^2+y^2 \leq 10} dxdy = 3(10\pi) = \boxed{30\pi}$$